Index,[1] Volume 52, 1959

Absorption of DDT by boll weevil, 17
Acalymma vittata, 477
Acaropsis sp., 443
Acarus siro, 237, 514
Aceratagallia
  sanguinolenta, 1145
  uhleri, 429, 430, 431
Aceria
  fixus, 186
  sp., 186
Acerophagus notativentris, 74
Acetaldehyde, acetals, toxicity of to house flies, 954
Acetic acid, methyl ester, toxicity of to house flies, 954
Acetophenone, tests of repellency of, 286
Acrididae, 431
Acrobeles ctenocephalus, 861
Acrobeloides
  bütschli, 861
  minor, 861
  tricornis, 861
Acrylamide, toxicity of to house flies, 954
Acrylic acid esters, toxicity of to house flies, 954
Acrylonitrile, effect of on
  confused flour beetle, 1091
  Sitophilus oryza, 1091
Acrylonitrile, toxicity of to Khapra beetle, 312
Actia interrupta, 550
Activity, Biological, of Hercules AC-528 in rats and cockroaches, 20
Adelphocoris
  lineolatus, 1145
  rapidus, 1145
Aedes
  aegypti, 263, 268, 977, 1027
  sollicitans, 1050
  taeniorhynchus, 356, 977
  vexans, 1050
Agapostemon melliventris, 1002
Agathis
  lacticinctus, 1103
  sp., 550
Aggregation, measurement of in insects, 1180
Agonoderus lecontei, 365, 1212
Agria affinis, 550
Agrilus auxius, 255
Agroperina dubitans, 966
Agrotis
  gladaria, 1212
  venerabilis, 966, 1212
Alabama argillacea, 757, 975
Aldrin
  breakdown of in soils, 535
  effect of on rate of seeding wheat, 535
  effect of seed treatment with on wireworms, 509
  persistence of in soils, 124, 289
Aldrin, for control of
  Acarus siro, 237
  alfalfa weevil, 663
  apple maggot, 751
  army cutworm, 784
  Drosophila melanogaster, 1136
  Drosophila spp., 755
  Hippelates pusio, 68
  imported fire ant, 1
  larger cabinet beetle, 234
  meadow spittlebug, 663
  mushroom flies, 257
  narcissus bulb fly, 348
  pea aphid, 663
  Profenusa canadensis, 1218
  sheep ked, 380
  strawberry weevils, 1117
  tobacco thrips, 451
Aldrin, effect of on
  chinch bug, 759
  garden symphylid, 666
  Japanese beetle, 9
  lesser cornstalk borer, 63
  potato leafhopper, 908
  Psila rosae, 963
  soil nematodes, 861
  tobacco flea beetle, 753
Aldrin, residues of
  in cattle, 1206
  on sweet potatoes, 1119
Aldrin, toxicity of to
  Douglas-fir beetle, 485
  garden symphylid, 892
  khapra beetle, 312
  spruce budworm, 212
Alectoris graeca, 774
Aleurocanthus woglumi, 1099
Aleurocybotus sp., 9
Aleurodothrips fasciapennis, 577
Aleyrodes sp., 9
Alkylphosphonic acid esters, effectiveness of as insecticides, 1121
Allethrin
  as house fly spray, comparison with chrysanthemumic acid esters, 521
  effect on action of by piperonyl butoxide, 1168
Allethrin, for control of
  Acarus siro, 237
  larger cabinet beetle, 234
  Tyrophagus spp., 237
Allethrin, effect of on house flies as space spray, 704
Allethrin, toxicity of to
  house flies, 85, 733
  khapra beetle, 312
Allothrombium sp., 231
Alloxan, for control of beet leafhopper, 701
Alsophilus pometaria, 1103
Amathes c-nigrum, 1212
Amblymerus bruchophagi, 898
Amblyomma americanum, 512
Amblyseiopsis, n. sp. 189
Am. Cyanamid 5223, effect of on honey bees, 549
Am. Cyanamid 12008, for control of
  citrus whitefly, 1008
  cotton thrips, 35
  Myzus cerasi, 778
Am. Cyanamid 12008, effect of on garden symphylid, 666
Am. Cyanamid 12009, for control of
  citrus whitefly, 1008
  Myzus cerasi, 778
Am. Cyanamid 12880 (see also dimethoate)
Am. Cyanamid 12880, effect of on
  alfalfa weevil, 758
  pea aphid, 758
Am. Cyanamid 18133
  for control of *Conoderus falli*, 780
  toxity of to onion maggot, 683
Am. Cyanamid 18133, effect of on
  garden symphylid, 672
  Psila rosae, 963
Am. Cyanamid 18706, toxicity of to onion maggot, 683
Am. Cyanamid 18706, for control of
  apple aphid, 493
  cabbage looper, 1013
  Mexican bean beetle, 541
  pea aphid, 541
  rosy apple aphid, 493
Am. Cyanamid 18706, effect of on
  alfalfa weevil, 758
  cabbage aphid, 816
  cabbage looper, 816
  pea aphid, 758
  Pieris rapae, 816
Amphimallon majalis, 171, 744
Amphorophora
  lactucae, 422, 423 424
  rhododendri, 423
  rubitoxica, 423
  sonchi, 6, 440
  viciae, 424
  sp., 422
Anarhopus sydneyensis, 75
Anarsia lineatella, 340, 637
Aneristus ceroplastae, 596
Anisaldehydes, toxicity of to house flies, 954
Anisota senatoria, 460, 1015
Anopheles
  crucians, 1050
  punctipennis, 1050
  quadrimaculatus, 868, 977, 1027, 1038
Anthonomus grandis, 134, 310, 346, 390, 403, 453, 466, 555, 592, 603, 710, 755, 757, 807, 840, 975, 985, 1010, 1018, 1042, 1068, 1099, 1138
Anthonomus grandis thurberiae, 603
Anthrenus flavipes, 382, 525
Anuraphis
  bakeri, 422
  helichrysi, 6, 422
  roseus, 377, 499
  tulipae, 6, 423
  spp., 422
Ancystus
  agilis, 1103
  sp., poss. *agilis*, 231
Aonidiella aurantii, 74, 857
Apanteles
  californicus, 531
  circumscriptus, 1099
  fumiferanae, 550
Aphalara
  gutierreziae, 8
  suaedae, 8
Aphalaroida, sp., 9
Aphelenchoides spp., 861
Aphelenchus avenae, 861
Aphelinus
  chaonia, 536
  semiflavus, 136, 142, 154, 537
  sp., 536
Aphidencyrtus
  aphidivorus, 537
  sp., 536
Aphidius sp., 429
Aphidophagous insects, economic effect of on citrus in China, 536
Aphis
  abbreviata, 424

[1] Compiled by Richard H. Foote

apii, 1057
cardui, 423
circii-acanthoidis, 424
craccae, 424
fabae, 6, 304, 422, 423, 424, 1057
gossypii, 6, 423, 424, 440, 477, 536, 632, 757, 807, 1057
helianthi, 6, 423
helichrysi, 424
medicaginis, 6, 440
mordwilkiana, 424
nerii, 6
pomi, 493
pseudohederae, 6
spiraecola, 6, 536
subterranea, 424
tetrapteralis, 6
urticata, 424
viciae, 424
Aphytis
chrysomphale, 577
lepidosaphes, 1025
mytilaspidis, 1103
Apis mellifera, 286, 399, 461, 547, 549, 572, 969, 1002, 1024
Aplomyia caesar, 550
Aramite, for control of
mites, 434
mites in insect cultures, 443
Tetranychus cinnabarinus, 760
two-spotted spider mite, 1026
Aramite, effect of on
alfalfa mites, 991
European red mite, 871
seed germination, 355
Araneida, 429, 430, 431
Arasan, effect of on yields of red clover, 1145
Argas persicus, 1195
L-Arginine, for control of beet leafhopper, 701
Armigeres subalbatus, 490
Arochlor 5460, effect of on larger cabinet beetle, 234
Aspergillus
amstelodami, 514
candidus, 514
flavus, 514
glaucus, 514
ochraceus, 514
repens, 514
restrictus, 514
ruber, 514
Aspidiotus perniciosus, 373
Aspidiotiphagus
citrinus, 577, 1026
lounsburyi, 577, 1026
Aspidiotus destructor, 1099
Astichus sp., 895
Attagenus piceus, 234, 382, 457, 525
Attractants
for *Culex tarsalis*, studies on, 496
for gypsy moth, stability of, 82
for *Hippelates pusio*, studies on, 496
for house flies, isolation of, 981
methods of testing of, 496
Augochlorella pomoniella, 1002
Autogeny in *Aedes taeniorhynchus*, 356

Bacillus
anthracis, 506
cereus, 506
entomodicus var. *entomodicus*, 506
entomodicus var. *subtoxicus*, 506
sotto, 798
Bacillus thuringiensis, 1226
var. *alesto*, 506
"Anduze" strain, 506
var. *sotto*, 506
var. *thuringiensis*, 506
Bacteria isolated from fly colony watering pads, 542
Baits
sprays for walnut husk fly control, 279
toxicity of to house flies, 345
Balaustium
aonidophagus, 189
sp., 189, 231
Barthrin and isomers as affected by piperonyl butoxide, 1168
Bayer 21/199 (see also Co-Ral)
effect of on goat lice, 980
metabolism of in cattle, 214
toxicity of to onion maggot, 683
Bayer 21/199, for control of
cattle grubs, 425, 512
sheep ked, 380, 539
Bayer, 21/199, mode of action in
house fly, 643
mouse, 660
Bayer 21/199, resistance to
of *Culicoides furens*, 352
of house flies, 308
Bayer 17450, for control of
Acarus siro, 237
larger cabinet beetle, 234
Bayer 18935, for control of
Acarus siro, 237
larger cabinet beetle, 234
Bayer 19596, for control of
Acarus siro, 237
larger cabinet beetle, 234
Bayer 19639
determination of residues of, 76
residual action of, 304
to prevent grasshopper damage, 37
Bayer 19639, for control of
cabbage aphid, 1045
cotton thrips, 35
Bayer 19639, effect of on
Japanese beetle, 9
Lygus bugs, 461
Bayer 22408, for control of winter tick, 1220
Bayer 25141
chemistry and mode of action of, 94
contact and systemic properties of, 99
Bayer 25141, effect of on
Brevicoryne brassicae, 99
house fly, 99
Laphygma frugiperda, 99
Metatetranychus citri, 99
Periplaneta americana, 99
Tribolium confusum, 99
Bayer 29493 (see Bayer S-1752)
Bayer S-1752, effect of on
Culex pipiens quinquefasciatus, 1096
house fly, 1096
Rhodnius prolixus, 1096
Bean yellow mosaic virus, transmission of, 260
Beauveria bassiama, 508
Bemisia sp., 9
Benzaldehyde, acetals, toxicity of to house flies, 954
Benzenes, toxicity of to house flies, 954
Benzyl benzoate, repellency tests of, 286
Berlese funnel
for collecting pink bollworm, 1010
modified, for sampling arthropod populations, 1085
BHC
effect of on yields of red clover, 1145
persistence of in turf soils, 289
BHC, for control of
cotton insects, 1068
cotton thrips, 35
Drosophila spp., 755
Hippelates pusio, 68
insects boring in pine logs, 173
spotted alfalfa aphid, 642
BHC, effect of on
black flies, 699
plum curculio, 921
BHC, toxicity of to
convergent lady beetle, 616
European pine shoot moth, 588
Orius insidiosus, 616
pink bollworm, 301
spruce budworm, 212
Bioassays, of residues of organophosphorus insecticides, 840
Biological control
basis of, 784
of nigra scale in California, 596
past, present and future, 1099
programs in, 1106
of spotted alfalfa aphid, 142
Biology of
alfalfa weevil, 939
Aphelinus chaonia, 536
Aphelinus semiflavus, 154
aphids, flying, 440
Asiatic oak weevil, 177
bee visitors to saguaro, 1002
beet leafhopper, 470
Black Hills beetle, 1199
black flies in Maryland, 629
Blaniulus guttulatus, 1021
boll weevil
diapause collecting methods, 346
feeding, oviposition preferences, 390
hibernation, 310
longevity, 453
in Louisiana, 596
on winecup, 755
cattle lice, 79
Centrinaspis capillatus, 730
Chrysops flavida, 884
clover root borer, 1143
Collops vittatus, 159
cotton rat in Dallas, Tex., 770
Crambus dorsipunctellus, 966
plumbifimbriellus, 966
trisectus, 966
Crymodes devastator, 966
Culex tritaeniorhynchus, 490
Dioryctria abietella, 561
Estigmene acrea, 1109
European corn borer
egg masses, 295
pupation vs. temperature, 409
European red mite, 871
fabric pests, attraction of stains to, 382
forest tent caterpillar, 1180
grasshoppers
in cultural control, 336
food habits of, 342
greenbug, honeydew deposition, 368
Gnorimoschema gudmanella, 117
Haplothrips clarisetis, 786
honey bee, 969
house flies,
effect of chemicals on, 61
mating, 236
Hylemya fugax, 503
imported fire ant
effect of cold on, 347
food habits, 455
khapra beetle, 312
lesser peach tree borer, 634
mites
on alfalfa, 991
on Texas grapefruit, 228
on Texas grapefruit leaves, 725
Musca autumnalis, 1030
Nantucket pine moth, 1180

Nepticula juglandifoliella, 690
Oencyrtus kuwanae, 764
peach twig borer larvae, 340
Philaenus leucophthalmus, 904
pink bollworm, 305
Pleurophorus caesus, 1215
Praon palitans, 154
predaceous earwig, 247
predators of spotted alfalfa aphid, 159
Pseudocneorhinus bifasciatus, 586
Pseudococcus citriculus, 880
purple scale, 1025
red-pine sawfly, 1180
spotted alfalfa aphid, 249, 368, 1023
spotted lady beetle, 843
spruce budworm, 1180
strawberry root weevils, 1117
Trioxys utilis, 154
Trogoderma spp., 830
Typhlodromus fallacies, 871
vetch bruchid, 955
Biotypes of pea aphid in So. Quebec, 293
Blacklight as attractant for European chafer, 744
Blaniulus guttulatus, 1021
Blattella germanica, 448, 544, 1063
Blissus leucopterus, 759
Blood, rapid collection of in insects, 538
Boleodorus thylactus, 861
Boliden, effect of on drywood termite, 1211
Bombus sonorus, 1002
Bordeaux 5-5-50, effect of on
Entomophthora spp., 28
Boron trioxide, toxicity of to mosquito larvae, 1027
Bovicola
bovis, 79
caprae, 980
limbatus, 980
Brachyrinus
ovatus, 3, 1117
rugostriatus, 1117
sulcatus, 3, 1117
Bracon politiventris, 550
Bradysia fenestralis, 257
Brevicoryne brassicae, 6, 99, 422, 423, 424, 477, 816, 1045, 1057
Brevipalpus
aeoloides, 186
australis, 231
californicus, 186
cardinalis, 186
lewisi, 186, 936
phoenicis, 231
spp., 725
Bryobia
drummondi, 186
praetiosa, 186, 377
Bruchus brachialis, 955
Bucculatrix gossypiella, 175
Bulan resistance to of Dilan-resistant house flies, 997
Burettes, modified- for topical application of insecticides, 683
Butylacetanilide, test of repellency of, 286

Caenurgina
crassiuscula, 1212
erechtea, 966
Cages for rearing, 773, 1227
Calcium arsenate, effect of on
bollweevil, 1042
cotton pests, 975
grey garden slug, 360
Calliephialtes comstockii, 768
Callitroga hominivorax, 1006, 1202, 1217
Camphene
repellency of to pea aphid, 106
tests of repellency of, 286
Camphor, tests of repellency of, 286
Camponotus pennsylvanicus, 530
Capitophorus
braggi, 422
ribis, 424
spp., 7, 422
Captan, effect of on
apple insect, 826
Entomophthora spp., 28
sorghum stands, 365
Carbamates, effect of on DDT- and malathion-resistant house flies, 178
Carbon dioxide, effect of repellents on evolution of from humans, 1157
Carbon disulfide, effect of on
Amphimallon majalis, 171
confused flour beetle, 1091
Sitophilus oryza, 1091
Carbon tetrachloride, effect of on
confused flour beetle, 1091
Sitophilus oryza, 1091
Carbowax 6000, as diluent for seed treatment chemicals, 509
Carpocapsa pomonella, 819, 1099, 1103
Carpophilus
lugubris, 640
spp., 972
Carulaspis visci, 1099
Cathartus quadricollis, 612
Cavariella
aegopodii, 1057
pastinacae, 424
CCl_4:CS_2
toxicity of to
Oryzaephilus surinamensis 942
rice weevil 942
Tribolium confusum 942
use of for tests of fumigants bioassays 942
Celcure, effect of on drywood termites, 1211
Cerasan, to increase stands of sorghum, 365
Ceratitis capitata, 1099
Ceratocystis ulmi, 902
Ceromasia auricaudata, 550
Cerosipha epilobiina, 424
Chaetocnema spp., 972
Chaitophorus
betulinus, 424
populellus, 7
viminalis, 7
Charcoal, activated, in control of spotted alfalfa aphid, 654
Chelonus phthorimaeae, 117
Chemonite effect of on drywood termite, 1211
Chermes piceae, 828
Chilocorus stigma, 577
Chiloplacus symmetricus, 861
Chionodes psiloptera, 966
Chipman R-6199
effect of on *Lygus* bugs, 461
prevention of Dutch elm disease transmission by, 902
resistance of *Tetranychus cinnabarinus* to, 760
Chipman R-6200, effect of on *Bucculatrix gossypiella*, 175
Chloranil, effect of on germination and seedling growth of wheat, 650
Chlorbenside (see also Mitox), for control of two-spotted spider mite, 342
Chlordane
effect of on yields of red clover, 1145
influence of on southern lyctus beetle, 357
persistence of in turf soils, 289
protection of stored woolens, 457
residues of on and in lemon, 42
Chlordane, for control of
Acarus siro, 237
cotton thrips, 35
Drosophila melanogaster, 1136
Drosophila spp., 755
Hippelates pusio, 68
imported fire ant, 1
larger cabinet beetle, 234
strawberry weevils, 1117
Chlordane, effect of on
Japanese beetle, 9
tobacco flea beetle, 753
Chlordane, resistance to of
brown dog tick, 332
Culicoides furens, 352
German cockroach, 544
Chlordane, toxicity of to
Douglas-fir beetle, 485
European pine shoot moth, 588
house flies, 733
Chlorobenzilate, effect of on
alfalfa mites, 991
European red mite, 871
seed germination, 355
Chlorobenzilate, for control of
mites in insect cultures, 443
Tetranychus marianae, 763
Chlorogenus callistrephi var. *vulgaris*, 686
Chloropicrin, effect of on
confused flour beetle, 1091
Sitophilus oryza, 1091
Chloropicrin, for control of garden symphylid, 678
Chlorthion, for control of
Acarus siro, 237
alfalfa weevil, 663
corn earworm, 1111
green peach aphid, 727
larger cabinet beetle, 234
meadow spittlebug, 663
pea aphid, 663
tobacco hornworm, 798
Chlorthion, effect of on
garden symphylid, 666, 672
Japanese beetle, 9
Chlorthion, toxicity of to
house flies, 733
khapra beetle, 312
Cholinesterase
activity of in cattle erythrocytes, 66
inhibition, to determine Bayer 19639 residue, 76
Choristoneura fumiferana, 212, 618, 1180
Chorizagrotis auxiliaris, 783
Chrysanthemumates, evaluation of as space sprays for house flies, 704
Chrysanthemumic acid esters, toxicity of to house flies 85, 521, 954
Chrysobothris femorata, 255
Chrysocharis
ainsliei, 895
parksi, 895
spp., 895
Chrysolophus pictus, 774
Chrysomphalus aonidum, 577
Chrysopa
aurata, 819
bimaculata, 577
californica, 596
cubana, 577
interupta, 577
plorabunda, 429, 430, 431, 577
rufilabris, 577
spp., 230, 577
Chrysops
flavida, 884
flavida reicherti, 884

Chytridiosus of Florida red scale, 577
Cimex lectularius, 448, 685
Cinnamaldehyde, acetal, toxicity of to house flies, 954
Circulifer tenellus, 470, 700
Cisseps (Scepsis) fulvicollis, 168
Clausenia purpurea, 880
Clemora smithii, 1099
Closterocerus
trifasciatus, 895
utahensis californicus, 690
Clover, red
effect of insecticides on yield of, 1145
injury to by *Sitonia* spp., 1155
Cnephasia virauregana, 1099
Cnephia metata, 629
Coccidae, 782
Coccinella novemnotata frauciscana, 148
Coccophagus
cowperi, 596
eritreaensis, 596
gurneyi, 880
isipingoensis, 596
pseudococci, 880
pulvinariae, 596
rusti, 72
scutellaris 74, 596
Coccus
hesperidum, 74, 596
pseudomognoliarum, 596
Coccygomimus pedolis, 550
COCS, compatibility of Thiodan with on potato foliage, 564
Coleomegilla maculata, 843
Colinus virginianus, 774
Collecting
apparatus using suction, 1085
technique for pink bollworm, 1010
method for shaft louse, 1215
Collops
quadrimaculatus, 431
vittatus, 159
Common names of insecticides,
changes in, 1032
coinage of, 1037
list, revised, 361
Comperiella bifasciata, 577
Conoderus falli, 772, 780
Conotrachelus nenuphar, 819, 921
Copper naphthenate
effect of on drywood termite, 1211
tests of with *Reticulitermes flavipes*, 533
Copper sulfate, compatibility of Thiodan with on potato foliage, 564
Co-Ral (see also Bayer 21/199), effect of on
horn fly, 1216
poultry ectoparasites, 1195
Psila rosae, 963
Co-Ral, for control of
beet leafhopper, 701
horn fly, 648
screw-worm, 1217
winter tick, 1220
Co-Ral, metabolism of in tissues of
cattle grub, 692
house fly, 692
mouse, 692
ox, 692
rat, 692
Corn, damage to by rice weevil, 1130
CP-7769, effect of on cotton pests, 975
CP 10502 effect of on *Lugus* bugs, 461
CP 10516 effect of on *Lygus* bugs, 461
CPD 8305, effect of on harlequin cabbage bug, 777
Crambus
bonifatellus, 966
dorsipunctellus, 966
leachellus cypridalis, 966
pascuellus, 966
plumbifimbriellus, 966
topiarius, 966
trisectus, 966
vulgivagellus, 966
Creosote, tests of with *Reticulitermes flavipes*, 533
Crymodes devastator, 966, 1212
Cryptogonus orbiculus, 536
Cryptolaemus montrouzieri, 1099
Cryptopeltis notata, 1219
Ctenocephalides felis, 770
Ctenucha virginica, 168
Culex
pipiene, 259, 490
quinquefasciatus, 1027, 1096
restuans, 1050
salinarius, 1050
stigmatosoma, 259
tarsalis, 259, 496
territans, 259, 1050
tritaeniorhynchus, 490
Culicoides furens, 352
Culiseta incidens, 259
Culture, of
screw-worm larvae, 1006
symbionts of confused flour beetle, 756
Cultural control of lesser cornstalk borer in California, 63
Cyclethrin, for control of *Acarus siro*, 237
Cyclohexylcyclohexanol, tests of repellency of, 286
Cycloneda sanguinea, 577
Cylisticus convexus, 621
Cyprex—(see Am. Cyanamid No. 5223)
Cyprinodontidae, 696
Cyrtepistomus castaneus, 177
Cyrtorhinus mundulus, 74
CZC, effect of on drywood termite, 1211

Dactynotus cichoricola, 424
Dacus dorsalis, 1099
Daphnia pulex, 1015
Dargida procincta, 966
Dasyneura gleditschiae, 712
DBC, effect of on house fly biology, 61
DBCP, effect of on garden symphylid, 666
for control of garden symphylid, 678
D-D mixture, for control of
Conoderus falli, 780
garden symphylid, 678
DDA, toxicity of to boll weevil, 19
DDD (see also TDE), toxicity of to khapra beetle, 312
DDE
conversion to of DDT by house flies, 19, 528
DDT
action of on resistant boll weevils, 403
analogs, as DDT synergists, 626
toxicity of to house flies, 626
biotic effect of on aphids and mites, 163
conversion to DDE by house flies, 528
evaluation of in lettuce yellows prevention, 686
fate of in larvicide suspension, 1038
injury to red fescues by, 176
metabolism of by boll weevil, 17
persistence of in soils, 124
protection of store woolens, 457
releasing neuroactive agent in cockroach, 1070
reproduction of house flies following treatment by, 765
resistance of *Culicoides furens* to, 352
resistance of house flies to, 308
resistance of European house flies to, 419
seasonal tolerance of potato flea beetle to, 519
DDT, for control of
Acarus siro, 237
arthropod disease vectors, 1050
alfalfa weevil, 940
bronze birch borer, 255
cabbage caterpillars, 221
cabbage looper, 1013
corn earworm, 338, 1111
cotton insects, 461, 1068
Drosophila spp., 755
European red mite, 936
fall armyworm, 838
flat headed apple tree borer, 255
Hippelates pusio, 68
house fly, 776
larger cabinet beetle, 234
lesser cornstalk borer, 329
lesser peach twig borer, 637
Linognathus vituli, 782
Lygus bugs, 244
mosquito larvae, 259
Pyrausta nubilalis, 49
sheep ked, 380
Stegasta basqueella, 468
tobacco thrips, 451
western peach tree borer, 804
DDT, effect of on
alfalfa mites, 991
American cockroach, 1070
beet armyworm, 877
black flies, 629
Bucculatrix gossypiella, 175
cabbage looper, 877
cotton pests, 975, 985
Entomophthora spp., 28
flour beetles, 1184
goat lice, 980
growth and fruiting of cotton, 396
herlequin cabbage bug, 777
imported cabbageworm, 877
lepidopterous larvae, 1115
Japanese beetle, 9
Lygus bugs, 461
mosquito larvae, 977
plum curculio, 921
potato leafhopper, 908
Psila rosae, 963
soil nematodes, 861
sugarcane borer, 821
sweetcorn pests, 972
tobacco flea beetle, 753
yields of red clover, 1145
DDT, residues of on
apple, 1053
peach, 109
sweet potato, 1119
DDT, toxicity of to
bark beetles, 323
convergent lady beetle, 616
Douglas-fir beetle, 485
European pine shoot moth, 588
house flies, 733
khapra beetle, 312
onion maggot, 683
Orius insidiosus, 616
pink bollworm, 301
DDT-dehydrochlorinase, inheritance of by house flies, 931
DDVP
dispenser for, 775
relation of structure of to activity, 270
toxic action of on house flies, 44
DDVP, for control of
alfalfa weevil, 663
Drosophila spp., 755
meadow spittlebug, 663
pea aphid, 663

Tendipes plumosus, 331
tobacco hornworm, 798
DDVP, effect of on
fathead minnows, 331
garden symphylid, 666
Japanese beetle, 9
Delnav (see also Hercules AC-528), for control of
citrus whitefly, 1008
tobacco hornworm, 798
winter tick, 1220
Delnav, effect of on
apple insects, 826
cabbage aphid, 816
cabbage looper 816
desert spider mite, 975
garden symphylid, 666, 672
goat lice, 980
horn fly, 1216
Pieris rapae, 816
potato leafhopper, 908
Delphene
effect of on poultry ectoparasites, 1195
for control of horn flies, 762
Demeton
effect of soils on uptake of, 1160
for prevention of lettuce yellows, 686
residues of on various vegetable, 741
resistance of *Tetranychus cinnabarinus* to, 760
Demeton, for control of
alfalfa aphid, 663
alfalfa weevil, 940
apple aphid, 493
beet leafhopper, 701
beet yellows virus, 910
citrus whitefly, 1008
cotton thirps, 35
green peach aphid, 727, 910
lettuce seed stem aphid, 411
meadow spittlebug, 663
mites, 432
Myzus cerasi, 778
pea aphid, 663
rosy apple aphid, 493
spotted alfalfa aphid, 714
Tendipes plumosus, 331
Tetranychus cinnabarinus, 760
Tetranychus marianae, 763
tomato suckfly, 1219
Demeton effect of on
alfalfa mites, 991
apple insects, 819
cotton pests, 975
Entomophthora spp., 28
European red mite, 871
fathead minnows, 331
harlequin cabbage bug, 777
insects in alfalfa, 428
Japanese beetle, 9
Lygus bugs, 461
Demethon thiolisomer, for control of cabbage aphid, 1045
Dendroctonus
brevicomis, 323
monticolae, 323
ponderosae, 1199
pseudotsugae, 482, 485
Density, relationship of to effect of insecticides, 1184
insecticides, 1184
Dermacentor
albipictus, 1220
variabilis, 770
Deroceras reticulatum, 360
Derostenus
arizonensis, 895
variipes, 895
Derris, toxicity of to Mexican bean beetle, 1017
Diabrotica
longicornis, 1212
undecimpunctata howardi, 477
Dialeurodes citri, 1008
Diapause
of boll weevil in Louisiana, 603
of boll weevil, effect of insecticides on, 1042
Diatraea
saccharalis, 821
spp., 1099
Diaulinopsis callichroma, 895
Diazinon
biotic effect of on pecan aphids and mites, 163
protection of stored woolens, 457
reproduction of house flies following treatment by, 765
resistance of house flies to, 308
Diazinon, effect of on
alfalfa mites, 991
cabbage aphid, 816
cabbage looper, 816
fathead minnow, 331
garden symphylid, 666
harlequin cabbage bug, 777
house fly, 704
Japanese beetle, 9
Lygus bugs, 461
Pieris rapae, 816
Psila rosae, 963
sap beetles in corn, 637
Diazinon, for control of
Acarus siro, 237
alfalfa weevil, 590, 663
apple aphid, 493
beet leafhopper, 701
black carpet beetle, 234
Conoderus falli, 780
Corn earworm, 1111
Drosophila melanogaster, 1136
Drosophila spp., 755
larger cabinet beetle, 234
lesser peach twig borer, 637
meadow spittlebug, 663
Mexican bean beetle, 541
pea aphid, 541, 663
potato aphid, 355
rosy apple aphid, 493
sheep ked, 380
spotted alfalfa aphid, 654
Tendipes plumosus, 331
Tetranychus marianae, 763
thrips on spinach, 540
tomato suckfly, 1219
Tyrophagus spp., 237
Diazinon, toxicity of to
resistant & non-resistant house flies, 719, 733
onion maggot, 683
plum curculio, 921
Dibrom, effect of on
alfalfa weevil, 758
cabbage aphid, 816
cabbage looper, 816
harlequin cabbage bug, 777
Lygus bugs, 461
pea aphid, 758
Pieris rapae, 816
potato leafhopper, 908
poultry ectoparasites, 1195
Dibrom effectiveness on vegetable crops, 477
Dibrom, for control of
alfalfa weevil, 590
bagworms, 353
cabbage looper, 1013
Drosophila spp., 755
lesser peach twig borer, 637
Mexican bean beetle, 541
pea aphid, 541
potato aphid, 355
thrips on spinach, 540
tomato suckfly, 1219
Dibrom, toxicity of to western cherry fruit fly, 769
Dibutyl adipate, tests of repellency of, 286
Dicapthon, effect of on
cotton pests, 975
desert spider mite, 975
horn fly, 1216
Psila rosae, 963
Dicapthon, for control of
alfalfa weevil, 663
Drosophila melanogaster, 1136
meadow spittlebug, 663
Mexican bean beetle, 541
pea aphid, 663
sheep ked, 380, 539
thrips on spinach, 540
Dicladocerus sp., 531
Didasia
opuntiae, 1002
rinconis, 1002
sp., 1002
Dieldrin, effect of on
alfalfa mites, 991
apple insects, 826
chinch bug, 759
cotton insects, 975
growth and fruiting of cotton, 396
honey bees, 549
Japanese beetle, 9
lesser cornstalk borer, 63
mosquito larvae, 977
potato insects, 564
potato leafhopper, 908
rate of seeding wheat, 535
seed treatment with on wireworms, 509
sweet corn insects, 972
Dieldrin for control of
Acarus siro, 237
alfalfa weevil, 663, 939
apple maggot, 751
army cutworm, 785
Centrinaspis capillatus, 730
corn earworm, 1111
cotton pests, 466
cotton thrips, 35
Drosophila melanogaster, 1136
Drosophila spp., 755
imported fire ant, 1
larger cabinet beetle, 234
meadow spittlebug, 663
onion maggot, 851
pea aphid, 663
Profenusa canadensis, 1218
rice water weevil, 14
sheep ked, 380
Stegasta basqueella, 468
thief ants, 365
tobacco thrips on peanuts, 451
western peach tree borer, 804
Dieldrin
influence of on southern lyctus beetle, 357
persistence of in turf soils, 289
for protection of stored woolens, 457
seasonal tolerance of potato flea beetle to, 519
susceptibility of European house flies to, 419
Dieldrin, residues of
in cattle, 1206
on and in lemons, 42
on sweet potatoes, 1119

Dieldrin, resistance to of
Culicoides furens, 352
house flies, 308
pear psylla, 530
Dieldrin, toxicity of to
convergent lady beetle, 616
Douglas-fir beetle, 485
house flies, 733
onion maggot, 683
Orius insidiosus, 616
pink bollworm, 301
plum curculion, 921
spruce budworm, 212
western bark beetle, 323
Diet for American cockroach containing gibberellic acid, 1214
Diet for boll weevil
feeding and oviposition on, 1138
inositol indispensable for, 1018
larvae, 710
Diethyltoluamide,
action of on mosquitoes, 263, 268
effect of on biting flies of cattle, 866
effect of on evolution of CO_2 and moisture in humans, 1157
fate of in guinea pigs, 928
Dilan
resistance to in house fllies, 997
toxicity of to house flies, 733
Dilan, for control of
alfalfa weevil, 663
house fly, 776
meadow spittlebug, 663
pea aphid, 663
Dimethoate, effect of on
cabbage aphid, 816
cabbage looper, 816
Pieris rapae, 816
Dimethoate, for control of
apple aphid, 493
cabbage looper, 1013
cattle grubs, 1004
green peach aphid, 727
Mexican bean beetle, 541
pea aphid, 541
rosy apple aphid, 493
tomato suckfly, 1219
Dimethoate (see also Am. Cyanamid 12880), metabolism of in cattle, 1190
Dimethoate, toxicity of to
European pine shoot moth, 588
onion maggot, 683
Dimethyl disulfide, for control of garden symphylid, 678
Dimethyl phthalate
action of on mosquitoes, 263
effect of on evolution of Co_2 and moisture from humans, 1157
Dinitrocresol, toxicity of to western bark beetles, 323
Dioryctria
abietella, 561
amatella, 561
Dipterex
acceptability and toxity of to house flies, 345
conversion of to DDVP, 44
dispenser for, 775
resistance of house flies to, 308
Dipterex, effect of on
Estigmene acrea, 1109
fathead minnow, 331
garden symphylid, 666
goat lice, 980
house flies, 704
Japanese beetle, 9
lepidopterous larvae, 1115
Lygus bugs, 461
Psila rosae, 963
Dipterex, for control of
Acarus siro, 237
beet leafhopper, 701
corn earworm, 1111
Drosophila melanogaster, 1136
larger cabinet beetle, 234
narcissus bulb fly, 348
Pyrausta nubilalis, 49
sheep ked, 539
strawberry weevil, 3
Tendipes plumosus, 331
tobacco hornworm, 798
walnut husk fly, 279
Dipterex, toxicity of to
house fly larvae, 44, 719
plum curculio, 921
Distribution
of alfalfa weevil, 939
of *Trogoderma* spp. in California, 830
Di-Syston
effect of on *Psila rosae*, 963
effect of temperature and plant species on, 435
Dis-Syston, for control of
alfalfa weevil, 663
apple aphid, 493
citrus whitefly, 1008
Drosophila spp., 755
meadow spittlebug, 663
pea aphid, 663
rosy apple aphid, 493
spotted alfalfa aphid, 654, 714
Dithane D-14, compatibility with Thiodan on potato foliage, 564
Dithane D-78, effect of on Entomophthora spp., 28
Dithylenchus spp., 861
Diversinerus smithi, 72
DMC effect of on house fly biology, 61
DN 289 for control of mites, 432
Dorylaimus
bastiani, 861
carteri, 861
clavicaudatus, 861
monhystera, 861
obscurus, 861
pratensis, 861
sublabiatus, 861
spp., 861
Dow ET-14 (see also ronnel)
Dow ET-14, toxicity of to house fly larvae 719
Dow ET-15, for control of
Acarus siro, 237
larger cabinet beetle, 234
Dow ET-15, toxicity of to house fly larvae, 719
Dow ET-57 (see also ronnel), for control of
Acarus siro, 237
cattle grubs, 425
larger cabinet beetle, 234
thrips on spinach, 540
Tyrophagus spp., 237
Dow-ET-57 (also Korlan), Residual effect of on Japanese beetle, 9
Dowco 109, effect of on poultry ectoparasites, 1195
Dowco 109, for control of
cattle grubs, 169, 749, 1022
sheep keds, 539
DPCP, effect of on garden symphylid, 666
Drosophila
melanogaster, 172, 559, 755, 1136
spp., 767
Dusts
desiccation of drywood termites by, 190
physical properties of derris, 1017
prevention of damage by *Kalotermes minor* by, 208
Dutch elm disease, transmission of by bark beetles, 902
Dylox, effect of on
alfalfa mites, 991
alfalfa weevil, 758
harlequin cabbage bug, 777
painted lady, 1017
pea aphid, 758
potato leafhopper, 908
poultry ectoparasites, 1195
Dylox, for control of
alfalfa weevil, 590
thrips on spinach, 540
tomato suckfly, 1219

Echinolaelaps echidninus, 770
Egg (insect)
control of by phosphate oil sprays, 377
of corn earworm, estimates of abundance, 747
of *Drosophila* spp., in tomatoes, 767
of European corn borer, eaten by spotted lady beetle, 843
of gypsy moth, parasitized by *Oencyrtus kuwanae*, 764
of honey bee, defective through starvation, 166
of meadow spittlebug, 240
of *Prodenia ornithogalli*, survival on jet aircraft, 299
Eggs (poultry)
effect of chicken body louse on production of, 358
Elachertus cacaeciae, 553
Elasmophalpus lignosellus, 63, 329
Empoasca
fabae, 429, 430, 431, 477, 564, 908
sp., 1145
Encoptolophus pallidus subgracilis, 337
Endrin
action of on resistant boll weevils, 403
coverage of cotton by 2 sprayer types, 592
effect of seed treatment with on wireworms, 502
effect of on successive generations of boll weevils, 555
Endrin, effect of on
alfalfa insects, 991
arthropod populations in alfalfa, 428
cotton insects, 975
Estigmene acrea, 1109
harlequin cabbage bug, 777
Japanese beetle, 9
lepidopterous larvae, 1115
lesser cornstalk borer, 63
Lygus bugs, 461
pecan aphids and mites, 163
potato leafhopper, 908
sugarcane borer, 821
sweet corn insects, 972
Endrin, for control of
alfalfa weevil, 663
apple maggot, 751
army cutworm, 785
birdsfoot trefoil insects, 836
cabbage caterpillars, 221
corn earworm, 1111
cotton insects, 1068
Drosophila spp., 755
lesser cornstalk borer, 329
lesser peach twig borer, 637
meadow spittlebug, 663
onion maggot, 851
pea aphid, 663
Pyrausta nubilalis, 49
Stegasta basqueella, 468

tobacco hornworm, 798
western peach tree borer, 804
Endrin, resistance to of
Culicoides furens, 352
pear psylla, 530
Endrin, toxicity of to
convergent lady beetle, 616
Douglas-fir beetle, 485
onion maggot, 683
Orius insidiosus, 616
pink bollworm, 301
spruce budworm, 212
western bark beetles, 323
ENT-21170, for control of
Conoderus falli, 780
Entomophagous insects,
use of in pest control, 71, 532
Entomophthora
coronata, 28
exitialis, 28, 142
ignobilis, 28
obscura, 28
virulenta, 28, 30
sp., 351
Eotetranychus
frosti, 187
malvastris, 187
sexmaculatus, 187
talisiae, 187
weldoni, 187
yumensis, 187
Ephestia cautella, 888
Ephialtes ontario, 550
Epicauta pestifera, 477
Epilachna varivestis, 477, 541, 1017
Epitragodes tomentosus, 577
Epitrix
cucumeris, 519, 564
hirtipennis, 753
EPN, toxicity of to western bark beetle, 323
EPN, effect of on
fathead minnows, 331
garden symphylid, 666, 672
EPN, for control of
mites, 434
Tendipes plumosus, 331
Equipment
electric micro-dispenser, 1019
holders for spray cards, 1029
Erdalith, effect of on drywood termite, 1211
Eretmocerus serius, 1099
Eriophyes pyri, 183, 187
Erythrocytes of cattle, activity of, 66
Estigmene acrea, 966
Estimates of corn earworm egg abundance, 747
Ethion
effectiveness of on vegetable crops, 477
for overwintering mite and aphid eggs, 377
for scale control on dormant trees, 373
Ethion, effect of on
apple insects, 826
California red scale, 857
citrus red mite, 857
cotton pests, 975
harlequin cabbage bug, 777
Lygus bugs, 461
potato leafhopper, 908
Psila rosae, 963
Ethion, for control of
alfalfa weevil, 590, 663
apple aphid, 493
beet leafhopper, 701
Drosophila melanogaster, 1136
European red mite, 936
lesser peach twig borer, 637
meadow spittlebug, 663
pea aphid, 663
rosy apple aphid, 493
Tetranychus marianae, 763
thrips on spinach, 540
tobacco hornworm, 798
tomato suckfly, 1219
Ethion, toxicity of to
onion maggot, 683
western cherry fruit fly, 769
Ethnyl carbinol, effect of on house fly biology, 61
Ethyl hexanediol
action of on mosquitoes, 263
effect of on evolution of CO_2 and moisture in humans, 1157
Ethyl parathion, effect of on
Estigmene acrea, 1109
Ethylene dibromide, effect of on
confused flour beetle, 1091
garden symphylid, 666
Sitophilus oryza, 1091
Ethylene dibromide, for control of
Brachyrinus ovatus, 3
Brachyrinus sulcatus, 3
confused flour beetle, 1081
Conoderus falli, 780
garden symphylid, 678
khapra beetle, 312
Nemocestes (Geoderces) incomptus, 3
Sciopithes obscurus, 3
Ethylene dichloride, effect of on
confused flour beetle, 1091
Sitophilus oryza, 1091
Ethylene dichloride, for control of
western peach tree borer, 804
Ethylene chlorobromide, toxicity of to khapra beetle, 312
Eubadizon gracile, 550
Eucephalobus
oxyuroides, 861
striatus, 861
Eulype hastata, 351
Eupelmella vesicularis, 898
Eupelmus
allynii, 898
atropupureus, 898
microzonus, 898
Eurytoma pini, 768
Eutetranychus banksi, 228, 725
Euxoa
messoria, 966
ochrogaster, 350, 966
septentrionalis, 966
Exochomus marginipennis childreni, 577
Exochus sp., 550

Fallowing, for control of *Conoderus falli*, 772
Fannia
canicularis, 530
scalaris, 530
Feltia ducens, 966, 1212
Fenusa pusilla, 712
Ferbam, effect of on
apple insects, 826
Entomophthora spp., 28
Fertilization of sweet corn as affected by Japanese beetle, 475
Fertilizers, application of insecticides with, 767
Fluorescence
blacklight for attracting European chafers, 744
for tracking honey bees in cotton, 572
Foliafume, for control of thrips on spinach, 540
Folidol, for control of cotton insects, 1068
Formaldehyde, for control of onion maggot, 851
Formalin, for control of woolen pests, 457
Frankliniella
exigua, 807
fusca, 451, 807
occidentalis, 35, 429, 430, 431, 540
tritici, 179, 807
sp., 757
Fumigation
effect of, on germination of
barley, 319
oats, 415
rice, 706
wheat, 51
effect of factors on reliability of bioassays of, 942
effect of soil to control weevil grubs in strawberries, 3
sorption of fumigants determined by gas chromatography, 1091
Fumigation, for control of
confused flour beetle, 1081
khapra beetle, 312
pea aphid, 1212
Fundulus ocellaris, 696
Fungi
attacking Florida red scale, 577
effect of insecticides and fungicides on, 28
germination of resting spores of, 30
of Lepidoptera, *B. thuringensis* for control of, 877
of stored grain, associated with mites, 514
transmission of Dutch elm disease, 902
Fungicides
effect of on fungi of spotted alfalfa aphid, 28
to increase sorghum stands, 365
Galeria melonella, 1010
Gargapeia solani, 477
General Chemical 3583, for control of corn earworm, 1111
General Chemical 3661, toxicity of to plum curculio, 921
Genetics, genetic control of queen honey bee size, 547
Genite, for control of European red mite, 871
Genite 923 for control of mites, 432
Geocoris
atricolor, 461
pallens, 461
punctipes, 461
punctipes sonoraenses, 159
sp., 431
Germination of
barley, effect of fumigation on, 319
oats, effect of fumigation on, 415
resting spores of *Entomophthora virulenta*, 30
rice, effect of fumigation on, 706
wheat, effect of fumigation on, 51
wheat, effect of Thimet and stickers on, 650
Gibberellic acid in diet of American cockroach, 1214
Glass, flashed opal, for detecting parasites, 782
L-(+)-glutamic acid, for control of beet leafhopper, 701
Glyodin, effect of an apple insects, 819, 826
Glypta fumiferanae, 550
Gnorimoschema
gudmanella, 117
operculella, 908

Goniocotes
bidentatus, 772
sp., 774
Goniodes sp., 774
Gonipterus? scutellatus, 1099
Graphiphora c-nigrum, 966
Grapholitha molesta, 109, 596
Grasses as hosts of English grain aphid and apple grain aphid, 994
Gravity, effect of on life cycle of vinegar fly, 559
Growth of corn affected by European corn borer, 567
Guthion
action of on resistant boll weevils, 403
bioassay of residues of, 840
effect of on growth and fruiting of cotton, 396
effect of on successive boll weevil generations, 555
estimation of residues of, 791
Guthion, effect of on
alfalfa mites, 991
alfalfa weevil, 758
apple insects, 826
Bucculatrix gossypiella, 175
cabbage aphid, 816
cabbage looper, 816
cotton insects, 975, 985
fathead minnows, 331
garden symphylid, 666
Japanese beetle, 9
lepidopterous larvae, 1115
Lygus bugs, 461
pea aphid, 758
Pieris rapae, 816
potato leafhopper, 908
Psila rosae, 963
Guthion, for control of
alfalfa weevil, 590, 663
apple aphid, 493
beet leafhopper, 701
cabbage caterpillars, 221
corn earworm, 1111
Drosophila spp., 755
European red mite, 936
green peach aphid, 727
lesser peach twig borer, 637
meadow spittlebug, 534, 663
pea aphid, 663
rosy apple aphid, 493
Stegasta basqueella, 468
Tendipes plumosus, 331
thrips on spinach, 540
tobacco hornworm, 798
tomato suckfly, 1219
western peach tree borer, 804
Guthion, toxicity of to
onion maggot, 683
pink bollworm, 301
plum curculio, 921
western cherry fruit fly, 769

Habrocytus medicaginis, 898
Hadrobregmus carinatus, 452
Haemobartonella muris, 543
Haemolaelaps glasgowi, 770
Halictus (*Chloralictus*) sp., 1002
Halticoptera aenea, 898
Haplothrips
clarisetis, 786
faurei, 819
Heliocotylenchus spp., 861
Heliothis
virescens, 767, 807, 1115
zea, 338, 466, 477, 757, 807, 972, 975, 985, 1068, 1111, 1115, 1212
Hemerobius pacificus, 596
Hemisarcoptes malus, 189, 577, 1103
Heptachlor, effect of on
alfalfa mites, 991
chinch bug, 759
cotton insects, 975
Estigmene acrea, 1109
Japanese beetle, 9
lesser cornstalk borer, 63
potato leafhopper, 908
Psila rosae, 963
soil nematodes, 861
sugarcane borer, 821
sweet corn insects, 972
tobacco flea beetle, 753
Heptachlor,
effect of seed treatment with on wireworms, 509
oral toxicity of to young calves, 1127
persistence of in soils, 124, 289
persistence of in soils, 124, 289
resistance of *Culicoides furens* to, 352
Heptachlor, for control of
alfalfa weevil, 590, 663, 940
apple maggot, 751
army cutworm, 785
birdsfoot trefoil insects, 836
Centrinaspis capillatus, 730
corn earworm, 1111
cotton thrips, 35
Drosophila melanogaster, 1136
Drosophila spp., 755
imported fire ant, 1
meadow spittlebug, 663
mosquito larvae, 259
narcissus bulb fly, 348
onion maggot, 851
pea aphid, 663
Pyrausta nubilalis, 49
sheep ked, 380
Stegasta basqueella, 468
strawberry weevils, 1117
thief ant, 365
tobacco thrips on peanuts, 451
Heptachlor, residues of
in cattle, 1206
on and in lemon, 42
on sweet potato, 1119
Heptachlor, toxicity of to
convergent lady beetle, 616
Douglas-fir beetle, 485
onion maggot, 683
Orius insidiosus, 616
pink bollworm 301
western bark beetles 323
Heptachlor epoxide, oral toxicity of to young calves, 1121
Hercules AC426-A, for control of
Acarus siro, 237
larger cabinet beetle, 234
Tyrophagus spp., 237
Hercules AC-528 (see also Delnav)
chromatography of, 20
for control of sheep ked, 539
mechanism of residue loss of, 111
metabolism of in rats and cockroaches, 20
Hermetia illucens, 917
Hexarthrum ulkei, 452
Hexyl mandelate, tests of repellency of, 286
L-(+)-histidine, for control of beet leafhopper, 701
Histiostoma sp., 187
Hippelotes
bishoppi, 496
pusio, 68, 496
Hippodamia
convergens, 142, 159, 429, 577, 616, 781
parenthesis, 147
quinquesignata punctulata, 147
sinuata, 147
Hirsutella sp., 577
Holders for spray assessment cards, 1029
Homalodisca insolita, 359
Hopkins Bioclimatic Law, egg development of meadow spittlebug, 240
Hoplopleura hirsuta, 770
Horogenes, sp., 550
Hosts
effect of on clover root borer, 1143
of *Trogoderma* spp., in California, 830
Humidity, effect of on attractants for house flies, 981
Hyaloides harti, 819
Hyalopterus
arundinis, 7, 424
atriplicis, 7, 422, 423
pruni, 377, 440
Hydrellia griseola, 348
Hydrocyanic acid, effect of on
confused flour beetle
germination of barley, 319
germination of oats, 415
germination of rice, 706
germination of wheat, 51
Sitophilus oryza, 1091
Hydrocyanic acid, toxicity of to khapra beetle, 312
Hylastinus obscurus, 658, 1143, 1166
Hylemya
antiqua, 683, 851
cilicrura, 430
fugax, 503
Hylurgopinus rufipes, 902
Hypera
meles, 286
nigrirostris, 286
postica, 590, 663, 758, 939
Hyphantria cunea, 1099
Hypoderma
bovis, 169, 488, 512, 524, 692, 749, 958, 1004, 1028, 1221, 1222
lineatum, 169, 425, 488, 512, 524, 749, 958, 1004, 1028, 1221, 1222
spp., 1022, 1223
Hypolin, for control of cattle grubs, 1223
Hypopteromalus, sp., 531
Hyposoter exiguae, 327
Hyssopus thymus, 768
Hysteroneura setariae, 7
Hystrichonychus gracilipes, 187

Icerya purchasi, 72, 1099
Indopol H-1500, for control of two-spotted spider mite, 1015
Inheritance of DDT-dehydrochlorinase by house flies, 931
Injury to
potatoes by *Blaniulus guttulatus*, 1021
red clover by *Sitona* spp., 1155
Inositol as dietary requirement for boll weevil, 1018
Insecticides
action of an feeding mosquitoes, 263
application of to onion maggot, 683
biotic effects of on aphids and mites, 163
common names of, 361
dusts, desiccation of termites with, 190
use in fumigant bioassays, 942
to increase sorghum stands, 365
locale of on citrus fruits, 723
metabolism of by house fly, 692
metabolism of by mouse, 692
metabolism of by ox, 692
metabolism of by rat, 692

penetration of to cockroach nerve cord, 813
persistence of in soils, 124, 289
physical properties of derris, 1017
for prevention of Dutch elm disease, 902
for prevention of grasshopper damage, 37
for prevention of lettuce yellows, 686
population changes in mites caused by, 991
for protection of stored woolens, 457
residual behavior of on and in citrus, 42
residual effect of on wheat, 703
residues of on lindane-dipped animals, 1031
residues of on several vegetables, 741
resistance of cattle and cattle grubs to, 1026
resistance of *Culicoides furens* to, 352
resistance of German cockroach to, 544
resistance of Orlando house fly colonies to, 308
resistance of pear psylla to, 530
resistance of *Tetranychus cinnabarinus* to, 760
rotation of for prevention of resistance, 432
screening of for house fly control, 954
as space sprays for fly control, 521, 704
susceptibility of European house flies to, 419
as synergists of DDT, 626
as systemics for cattle grub control, 425
Insecticides, effect of on
cattle, as aerosols, 866
citrus leaf photosynthesis, 88
fungi parasitic on spotted alfalfa aphid, 28
growth and fruiting of cotton, 396
honey bees, 549
photosynthesis of lemon leaves, 883
germination of barley, 319
germination of oats, 415
germination of rice, 706
germination of wheat, 51, 650
rate of seeding wheat, 535
reproduction of resistant house flies, 765
rice germination, 706
soil nematodes, 861
total arthropod populations in alfalfa, 428
wheat germination and seedling growth, 650
Ionization, effect of on penetration of insecticide, 812
Ischnus inquisitorius atricpes, 550
Isodrin
effect of on *Psila rosae*, 963
for control of onion maggot, 851
toxicity of to Douglas fir beetle, 485
toxicity of to western bark beetles, 323
Isolan
effect of on alfalfa mites, 991
effect of soils on uptake of, 1160
for control of citrus whitefly, 1008
toxicity of to house flies, 733
Itoplectus
behrensii, 531
conquisitor, 768
4-cingulatus, 550
viduata, 550

Kalotermes minor, 208, 1211
Keltane, effect of on
apple insects, 826
desert spider mite, 975
European red mite, 871
seed germination, 355
Kelthane, for control of
beet leafhopper, 701
European red mite, 936
mites, 432
mites in insect cultures, 443
Tetranychus cinnabarinus, 760
Tetranychus marianae, 763
Kepone, effect of on
apple insects, 826
cabbage aphid, 816
cabbage looper, 816
Pieris rapae, 816
Psila rosae, 963
Kepone, for control of corn earworm, 1111
Kepone, toxicity of to plum curculio, 921
Korlan (see also ronnel)
effect of on poultry ectoparasites, 1195
for control of sheep ked , 380, 539
toxicity of to western cherry fruit fly, 769

Labidura riporia, 247
Lacinipolia
pensilis, 966
renigera, 1212
Laemophloeus ferrugineus, 131, 612
Lampetia equestris, 348
Laphygma
exigua, 327, 877
frugiperda, 99, 838, 972
Latex paint, for use with insecticides, 347
Lead arsenate, for control of apple insects, 826
lesser peach twig borer, 637
Lecaniobius cockerelli, 596
Lecanium
corni, 373, 596
pruinosum, 373
Lepidoptera associated with grasses, 966
Lepidosaphes
beckii, 1025
ulmi, 1103
Leptothrips mali, 819
Leptinotarsa decemlineata, 564
Leptus sp., 189
Lethane 384
effect of on flour beetles, 1184
toxicity of to house flies, 733
Leucania
farcta, 966
insueta heterodoxa, 966
luteopallens, 966
multilinea, 966
sp., 1212
Leuronota maculata, 9
Lime, compatibility with Thiodan on potato foliage, 564
Lindane
action of on resistant boll weevils, 403
breakdown of in soils, 118
crystals of for protection of stored woolens, 457, 525
effect of seed treatment with on wireworms, 509
in milk from sprayed cows, 349
persistence of in soils, 124
residues of in dipped sheep and goats, 1031
taste and odor in potatoes and red kidney beans, 473
Lindane, effect of on
black flies, 629
garden symphylid, 672
goat lice, 980
insects attacking Douglas-fir logs, 482
Japanese beetle, 9
Lygus bugs, 461
mosquito larvae, 977
soil nematodes, 861
Lindane, for control of
Acarus siro, 237
alfalfa weevil, 590, 663
birdsfoot trefoil insects, 836
black carpet beetle, 234
Drosophila spp., 755
larger cabinet beetle, 234
lettuce seed stem aphid, 411
Linognathus vituli, 782
meadow spittlebug, 534, 663
pea aphid, 663
Profenusa canadensis, 1218
sheep ked, 380
spotted alfalfa aphid, 654
strawberry weevils, 1117
thief ants, 365
tobacco thrips on peanuts, 451
Tyrophagus spp., 237
Lindane, resistance to of
brown dog tick, 332
Culicoides furens, 352
European house flies, 419
German cockroach, 544
house flies, 308
Lindane, toxicity of to
Douglas-fir beetle, 485
house flies, 733
khapra beetle, 312
western bark beetles, 323
Lindorus lophantae, 577
Linognathus
stenopsis, 980
vituli, 782
Liorhyssus hyalinus, 14, 242
Lipoplexis sp., 536
Liodontomerus
insuetus, 898
longfellowi, 898
perplexus, 898
Lipurus baculus, 772
Liriomyza pictella, 895
Lissorhoptrus oryzophilus, 767
Listrophonidae, 770
Lithocolletic messaniella, 1099
Lithophane sp., 1103
Loemophloeus minutus, 888
Lophortyx californicus, 774
Lyctus planicollis, 357
Lygocerus sp., 75
Lygus
elisus, 244
hesperus, 242, 244, 461
lineolaris, 286, 429, 430, 431, 1145, 1212
spp., 939
Lypha setifacies, 550

Macrocentrus ancylivorus, 596
Macrocheles muscae domesticae, 530
Macrosiphoniella
sanborni, 7
spp., 422
Macrosiphum
ambrosiae, 7, 440
barri, 7, 411, 423, 440
dirhodum, 442, 423
granarium, 7, 422, 423, 994
obscurus, 424
pisi, 7, 106, 260, 286, 293, 422, 423, 424, 429, 430, 431, 439, 541, 663, 758, 1057, 1160, 1212
rosae, 423
scoliopi, 422
solanifolii, 355, 422, 423, 424, 440, 477, 1057
tanaceti, 424
spp., 422

Macrosteles fascifrons, 430, 686
Madremyia saundersii, 550
Malacosoma
americana, 299
disstria, 1180
Malathion
acceptability and toxicity to of house flies, 345
bioassay of residues of, 840
biotic effect of on pecan aphids and mites, 163
effect of grain moisture content on, 131
effect of on vegetable crops, 477
lethal action of on house flies, 218
metabolites of, 1063
odor of in wheat, 1013
for prevention of lettuce yellows, 686
residues of in milk, 341
residues of on and in orange, 42
Malathion, effect of on
apple insects, 826
arthropod population in alfalfa, 428
black flies, 629
cabbage aphid, 816
cabbage looper, 816
California red scale, 857
chinch bug, 759
citrus red mite, 857
cotton pests, 975
Entomophthora spp., 28
fathead minnows, 331
house flies, 704
Japanese beetle, 9
lepidopterous larvae, 1115
Lygus bugs, 461
mosquito larvae, 977
Pieris rapae, 816
Psila rosae, 963
sap beetles in corn, 640
seed germination, 353
sweet corn pests, 972
Malathion, for control of
Acarus siro, 237
alfalfa weevil, 590, 663, 939
arthropod disease vectors, 1050
birdsfoot trefoil insects, 836
black carpet beetle, 234
bronze birch borer, 255
cabbage caterpillars, 221
cabbage looper, 1013
citrus whitefly, 1008
corn earworm, 338, 1111
cotton thrips, 35
Drosophila melanogaster, 1136
Drosophila spp., 755
flat headed apple tree borer, 255
horn flies, 762
larger cabinet beetle, 234
lettuce seed stem aphid, 411
lice of domestic game birds, 774
Linognathus vituli, 782
mange mites, 790
meadow spittlebug, 663
Mexican bean beetle, 541
mosquito larvae, 259
northern fowl mite, 13
onion maggot, 851
pea aphid, 663
pigeon lice, 772
potato aphid, 355
Profenusa canadensis, 1218
scale insects on dormant fruit trees, 373
sheep ked, 380, 539
Stegana basqueella, 468
Tendipes plumosus, 331
thrips on spinach, 540
tomato suckfly, 1219
Tyroglyphus sp., 237
walnut husk fly, 279
winter tick, 1220
Malathion, resistance to of
Culicoides furens, 352
European house flies, 419
German cockroach, 544
house flies, 308
Malathion, toxicity of to
American cockroach, 1063
European pine shoot moth, 588
garden symphylid, 892
German cockroach, 1063
house flies, 733, 1063
house fly larvae, 719
khapra beetle, 312
killifish, 696
mouse, 1063
pink bollworm, 301
plum curculio, 921
Marietta sp., 596
Mediolata spp., 577, 819
Megaselia
agarici, 257
sp., 1010
Melanocallis caryaefoliae, 163
Melanophila californica, 323
Melanoplus
bilituratus, 336
bivittatus, 37
differentialis, 336
Melanotrichus flavosparsus, 1212
Melanotus spp., 509
Melissodes sp., 1002
Melophagus ovinus, 380, 539
Menacanthus stramineus, 358, 1195
Menopon gallinae, 1215
Mercaphos, for control of larger cabinet beetle, 234
Metabolism
of Bayer 21/199 in cattle, 214
of DDT in boll weevil, 17
of dimethoate in cattle, 1190
of Hercules AC-528 components in rats and cockroaches, 20
of Prolan in Dilon-resistant house flies, 997
Metaboric acid, toxicity of to mosquito larvae, 1027
Metaldehyde, effect of on grey garden slug, 360
Metaphycus
helvolus, 74, 596
luteolus, 74, 596
stanleyi, 74
Meteorus
trachynotus, 550
sp., 531
dl-Methionine, for control of beet leafhopper, 701
Methoxychlor
for protection of stored woolens, 457
residues of in milk, 866
Methoxychlor, effect of on
apple insects, 826
biting flies of cattle, 866
goat lice, 980
horn flies, 1216
Japanese beetle, 9
Methoxychlor, for control of
Acarus siro, 237
alfalfa weevil, 590, 663
Centrinaspis capillatus, 730
Drosophila melanogaster, 1136
horn flies, 762
larger cabinet beetle, 234
meadow spittlebug, 663
pea aphid, 63
sheep ked, 380
tomato suckfly, 1219
Methoxychlor, toxicity of to
house flies, 733
plum curculio, 921
Methyl bromide, effect of on
confused flour beetle, 1091
germination of barley, 319
germination of oats, 415
germination of rice, 706
germination of wheat, 51
Sitophilus oryza, 1091
Methyl bromide, for control of
confused flour beetle, 1081
garden symphylid, 678
Methyl bromide, toxicity of to
khapra beetle, 312
Methyl carbamate, effect of on DDT and Malathion-resistant house flies, 118
Methyl cellulose, for control of spotted alfalfa aphid, 714
Methyl demeton, for control of citrus whitefly 1008
Methyl parathion
bioassay of residues of, 840
for control of *Tendipes plumosus*, 331
Methyl parathion, effect of on
boll weevil, 1042
chinch bug, 759
cotton pests, 975
Estigmene acrea, 1109
fathead minnow, 331
Methyl parathion, toxicity of to
convergent lady beetle, 616
onion maggot, 683
Orius insidiosus, 616
Methyl pentanone repellency of to pea aphid, 106
Methyl trithion, for control of tomato suckfly, 1219
Metopeurum fuscoviride, 424
Metopolophium occidentale, 424
Mice
action of Bayer 21/199 and piperonyl butoxide in, 660
control of mange mites on, 790
metabolism of Co-Ral in tissues of, 692
Microweisia coccidivora, 577
Migration of *Frankliniella tritici* through greenhouse ventilators, 179
Mirini, 895
Mites, grain storage fungi associated with, 514
Mitox (see also Chlorbenzide), for control of European red mite, 871
Moisture, effect of
on clover root borers, 658, 1166
repellents on evolution of in humans, 1157
Monarthropalpus buxi, 712
Monellia costalis, 7, 163
Monomorium pharaonis, 448
Mononchus parabrachyurus, 861
Monsanto CP-7769 effect of on *Bucculatrix gossypiella*, 175
Monsanto 8574, for control of corn earworm, 1111
Monsanto 10502, for control of corn earworm, 1111
Monsanto 10851, for control of corn earworm, 1111
Monsanto 11327, for control of corn earworm, 1111
Murgantia histionica, 777
Musca
autumnalis, 1030
domestica, 23, 44, 61, 85, 99, 218, 236, 308, 345, 419, 446, 448, 521, 542, 626, 643, 692, 704, 719, 733, 765, 775, 776,

779, 917, 931, 954, 981, 997, 1050, 1063, 1077, 1096, 1121, 1168
Mutation of *Bacillus thuringensis* to *B. anthrocis*, 506
Myiophagus sp., 577
Mylone, effect of on garden symphylid, 672
Myobia musculi, 790
Myocoptes musculinus, 790
Myzus
ascalonicus, 424
cerasi, 377, 778
circumflexus, 423, 424
ligustri, 424
ornatus, 424
persicae, 7, 249, 260, 422, 423, 424, 440, 536, 727, 781, 910, 1045, 1057
solani, 424
spp., 422

Nabis
alternatus, 429, 430, 431
ferus, 159
Naphthenic oils, influence of on plant transpiration, 334
Nasonovia ribicola, 424
Natural control of insect populations, 532
Musca domestica by *Hermetia illucens*, 917
Novadel, for control of lesser peach twig borer, 637
NC-13 relation of structure of to insecticidal activity, 270
Nemagon, for control of
Conoderus falli, 780
strawberry weevils, 3
Nemocestes (*Geoderces*) *incomptus*, 3
Nemorilla floralis, 550
Neotetranychus siccus, 187
Neodiprion nanulus, 600, 1180
Nephelodes emmedonia, 966, 1212
Nepticula juglandifoliella, 690
Neuroactive substance, release of by DDT in cockroach, 1070
Nicarbazin, for control of cattle grubs, 1222
Nodita floridana, 577
Nosema sp., 351
Numbers of insects, description of, 847
Nutrients, aphids on oats grown in, 624
Nysius raphanus, 429

Obituaries
Clair Cahill Alexander, 1033
Ernest Adna Back, 1229
Thomas R. Chamberlin, 118
John E. Dudley, Jr., 450
Charles Lewis Fluke, 787
Francis B. Foley, 1034
Freeman M. Fuller, Jr., 788
Frederick B. LaForge, 180
Harold S. McConnell, 180
Harris Braley Parkes, 362
Altus L. Quaintance, 182
Willis Everett Snow, 1229
Harley Frost Wilson, 788
Clell Burns Wisecup, 789
James Zetek, 1230
Odor, in malathion-treated wheat, 1013
Oencyrtus kuwanae, 764
Oil sprays, effect of on
California red scale, 857
citrus red mite, 857
photosynthesis, 88, 174
plant transpiration, 334
Oil sprays, for control of
overwintering mite and aphid eggs, 377
scale insects on dormant fruit trees, 373
Olfactometer, a modified model, 621
Oligonychus
coniferarum, 187
modestus, 187
platani, 187
pratensis, 187
stickneyi, 187
ununguis, 187
Olla
abdominalis, 147
abdominalis var *plagiata*, 577
Omotoma fumiferanae, 550
Oncopeltus fasciatus, 177
Onchophanes americanus, 550
1,2-D, for control of garden symphylid, 678
Operophtera brumata, 1103
Opius
suturalis, 895
spp., 895
Orchopeas leucopus, 770
Oregma lanigera, 536
Organophosphates, penetration of in cockroach, 812
Origilus sp., 117
Orius
insidiosus, 429, 430, 431, 616, 819
tristicolor, 159, 461
Ornithonyssus
bacoti, 770
sylviarum, 13, 170, 1195
Oryzaephilus surinamensis, 557, 612, 942
OS-1808, relation of structure of to insecticidal activity, 270
OS-1836, relation of structure of to insecticidal activity, 270
Osmose, tests with *Retieulitermes flavipes*, 533
Ovex, for control of
European red mite, 871
mites, 432
mites in insect cultures, 443
two-spotted spider mite, 342, 1026
Ovex, effect of on seed germination, 355
Ox, metabolism of Co-Ral in tissues of, 692
Oxya chinensis, 73

Pachycrepoideus vindemmiae, 530
Panagrolaimus subelongatus, 861
Panogen to increase stands of sorghum, 365
Panonychus
citri, 857
ulmi, 377, 432, 871, 936, 1103
Paradichlorobenzene
for control of western peach tree borer, 804
for protection of stored woolens, 457
repellency of to pea aphid, 106
tests of repellency of, 286
Paraffinic oils, influence of on plant transpiration, 334
Paraphlepsius irroratus, 1212
Parasites of
Fannia canicularis and *F. scalaris*, 530
Florida red scale, 577
gypsy moth eggs, 764
Laphygma exigua, 327
melon leaf miner, 895
Nepticula juglandifoliella, 690
Pseudococcus citriculus, 880
Rhyacionia buoliana, 768
spotted alfalfa aphid in California, 136, 142, 154
spruce budworm in Oregon, 550
Parasites, utilization of, 1099
Parasitus, sp., 189
Parathion, effect of on
alfalfa mites, 991
Entomophthora spp., 28
fathead minnows, 331
garden symphylid, 666, 672
harlequin cabbage bug, 777
Japanese beetle, 9
Lygus bugs, 461
mosquito larvae, 977
painted lady, 1017
potato insects, 564
Psila rosae, 963
red clover yields, 1145
sap beetles in corn, 640
sweet corn pests, 972
Parathion, for control of
alfalfa weevil, 590, 939
birdsfoot trefoil insects, 836
bronze birch borer, 255
cabbage caterpillars, 221
cabbage looper, 1013
citrus whitefly, 1008
Conoderus falli, 772, 780
Drosophila spp., 755
flat headed apple tree borer, 255
garden symphylid, 678
green peach aphid, 727
lesser peach twig borer, 637
lettuce seed stem aphid, 411
mites, 434
scale insects on dormant fruit trees, 373
Tendipes plumosus, 331
thrips on spinach, 540
tobacco hornworm, 798
tomato suckfly, 1219
Parathion
for prevention of lettuce yellows, 686
residues of on apple, 1053
Parathion, resistance to of
Culicoides furens, 352
house flies, 308
Tetranychus cinnabarinus, 760
Parathion, toxicity of to
convergent lady beetle, 616
garden symphylid, 892
house flies, 733
onion maggot, 683
Orius insidiosus, 616
pink bollworm, 301
plum curculio, 921
western cherry fruit fly, 769
Paratrioza cockerelli, 9
Parlatoria oleae, 373
Pectinophora gossypiella, 301, 305, 385, 466, 1010
Pediculus humanus humanus, 448
Penicillium sp., 514
Pentachlorphenol, tests of with *Reticulitermes flavipes*, 533
Pentalonia nigronervosa, 424
Pentamerismus sp. nr. *baptus*, 187
Perdita spp., 1002
Periplaneta americana, 99, 448, 538, 1063, 1070, 1214, 1227
Perkinsiella saccharicida, 74
Perthane, effect of on
alfalfa weevil, 663
cabbage aphid, 816
cabbage looper, 816
harlequin cabbage bug, 777
Japanese beetle, 9
larger cabinet beetle, 234
meadow spittlebug, 663
pea aphid, 663
Pieris rapae, 816
Psila rosae, 963
Perthane, for control of
Acarus siro, 237

cabbage looper, 1013
corn earworm, 1111
lesser peach twig borer, 637
potato aphid, 355
thrips on spinach, 540
tomato suckfly, 1219
Petroleum oils, effect of on
photosynthesis of lemon leaves, 883
Petrobia latens, 187
Phaeogenes
hariolus, 550
sp., 531
Phasianus
cholchicus, 774
colchicus mongolicus, 774
Phenol, acetates of, toxicity of to house flies, 954
Phenothiazine
for control of cattle grubs, 1223
toxicity of to house fly larvae, 719
Philaenus
leucophthalmus, 240, 534, 663, 836, 904
var. *fabricii*, 904
var. *fasciatus*, 904
var. *leucocephalus*, 904
var. *leucophthalmus*, 904
var. *marginellus*, 904
var. *pallidus*, 904
var. *spumarius*, 904
Phorate (see also Thimet), effect of soils on uptake of, 1160
Phorocera
erecta, 550
incrassata, 550
Phryxe pecosensis, 550
Phosdrin, effect of on
Anisota senatoria, 1015
chinch bug, 759
fathead minnows, 331
harlequin cabbage bug, 777
Japanese beetle, 9
lepidopterous larvae, 1115
painted lady, 1017
Tendipes plumosus, 331
Phosdrin, for control of
alfalfa weevil, 590, 663, 939
beet leafhopper, 701
birdsfoot trefoil insects, 836
Cabbage caterpillars, 221
corn earworm, 338, 1111
meadow spittlebug, 663
pea aphid, 663
thrips on spinach, 540
tomato suckfly, 1219
Phosdrin
effect of soils on uptake of, 1160
relation of structure of to activity, 270
toxicity of to plum curculio, 921
Phosphamidon, effect of on *Psila rosae*, 963
Phosphamidon, for control of
beat leafhopper, 701
Mexican bean beetle, 541
pea aphid, 541
Phosphoric acid, esters of, toxicity of to house flies, 954
Phostex, for control of
Acarus siro, 237
alfalfa weevil, 663
apple aphid, 493
Eriophyes pyri, 183
larger cabinet beetle, 234
meadow spittlebug, 663
overwintering mite and aphid eggs, 377
pea aphid, 663
rosy apple aphid, 493
scale insects on dormant fruit trees, 373
Photosynthesis, citrus, effect on of
water phase of oil spray, 174
oil type, deposit, and soaking, 88
petroleum oils, 883
Phyllocoptruta oleivora, 228, 725
Phyllotreta cruciferae, 477
Physiology, respiration rates of larval honey bees, 399
Phytocorus sp., 819
Phytodietus fumiferanae, 550
Phytomyza ilicis, 712
Pieris rapae, 221, 477, 816, 877
Pigeons, malathion for control of lice on, 772
Pimepales promelas, 331
Pineus sp., 828
Piperitone, toxicity of to house flies, 954
Piperonal, diethyl acetal, toxicity of to house flies, 954
Piperonyl butoxide, effect of on
Acarus siro, 237
action of barthrin and isomers, 1168
action of Bayer 21/199, 643, 660
black carpet beetle, 234
larger cabinet beetle, 234
Tyroglyphus sp., 237
Piperonyl butoxide
lethal action of with malathion on house flies, 218
protection of dried citrus pulp with, 224
residual behavior of on wheat, 703
Planococcus citri, 880
Plastics, bags for protection of woolens against insects, 457
Platyperigea extima, 966
Plectus spp., 861
Pleurophorus caesus, 1215
Plodia interpunctella, 557, 612, 888, 1226
Plutella maculipennis, 211
Plutella maculipennis, 211
Polistes exclamans, 798
Pollination of saguaro flowers by honey bee, 1002
Polybor 3, for control of house flies in caged layers, 446
Polybutenes, for control of two-spotted spider mite, 1015
Polygenis gwyni, 770
Polyhedral bodies in orange-striped oakworm, 460
Polynema sp., 895
Polyplax spinulosa, 543
Popillea japonica, 9, 289, 475, 771
Porthetria dispar, 83, 764
Potamyia flava, 1212
Praon palitans, 136, 142, 154
Pratylenchus
brachyurus, 861
minyus, 861
spp., 861
Predators
of Florida red scale, 577
of mites on Texas grapefruit, 229
of spotted alfalfa aphid, 159
utilization of, 1103
Prociphilus tessellatus, 353
Prodenia
eridania, 1115
ornithogalli, 299
Profenusa canadensis, 1218
Prolan, resistance to of Dilan-resistant house flies, 997
Pronematus
californicus, 189
ubiquitus, 231
Propionaldehyde, acetal of, toxicity of to house flies, 954
Propionic acid esters, toxicity of to house flies, 954
Propylene dichloride, for control of western peach tree borer, 804
Prosimulium
hirtipes, 629
magnum, 629
Prospaltella aurantii, 577, 1026
Protagrotis obscura, 966
Protein hydrolysates, bait sprays for control of walnut husk fly, 279
Protoleucania albilinea, 966
Protoparce
quinquemaculata, 761
sexta, 761, 798, 1115
Psallus seriatus, 757, 1068
Pseudaletia unipuncta, 1111, 1212
Pseudohomalopoda prima, 577
Pseudocneorhinus bifasciatus, 586
Pseudococcus
aonidum, 75
citriculus, 880
comstocki, 880
gahani, 880
maritimus, 74
Psila rosae, 963
Psilenchus
aberrans, 861
gracilis, 861
hilarulus, 861
Psorophora
ciliata, 1050
confinnis, 14, 1050
cyanescens, 1050
discolor, 14, 1050
Psylla pyricola, 530
Pulvinaria psidii, 1099
PVP-Iodine-10, effect of on garden symphylid, 666
Pyran derivative, toxicity of to house flies, 954
Pyrausta nubilalis, 49, 295, 409, 567, 843, 972, 1111, 1171, 1212
Pyrethrins,
compared with chrysanthemumic acid esters, 521
effect of action of by piperonyl butoxide, 1168
residual behavior of on wheat, 703
Pyrethrins, effect of on flour beetles, 1184
house flies, 704
Pyrethrins, for control of
Acarus siro, 237
larger cabinet beetle, 234
Tyroglyphus spp., 237
Pyrethrins, toxicity of to house flies, 55, 733
Pyrethrum for protection of dried citrus pulp, 224
wheat stored in ships holds, 557
Pyrexel for control of thrips on spinach, 540
Pyrocide 175, toxicity of to khapra beetle, 312
Pyrolan
effect of on DDT- and malathion-resistant house flies, 178
toxicity of to house flies, 733

Racks for vial storage, 1225
Radiation, effect of on *Anopheles quadrimaculatus*, 868
insects affecting man, 448
stored grain insects, electromagnetic, 888
Radioactive tracers in
Bayer 21/199 metabolic studies, 214
Disyston metabolic studies, 435

diethyltoluamide, 929
mosquito rearing water, 977
mosquito repellent, 268
Raphignathus sp. undescribed, 189
Rat, metabolism of Co-Ral in tissues of, 692
RE 4355, effect of on Japanese beetle, 9
Rearing of
bed bug, 685
boll weevil, 710
Dioryctria abietella, 561
mosquitoes, 977
screw-worm larvae, 1006
Trombicula akamushi, 824
Revurvaria
milleri, 531
starki, 847
Repellents
action of on mosquitoes, 263, 268
behavior of several insects toward, 286
effect of grain moisture content on, of malathion, 131
effect of on evolution of CO_2 and moisture from humans, 1157
response of pea aphid to, 106
Reproduction following treatment of resistant house flies, 765
Residues of
aldrin in cattle, 1026
Bayer 19639, 76, 304
Bayer 19639, detection of, 76
Bayer S-1752, against insects, 1096
chlordane, on and in lemons, 42
chlorinated hydrocarbons on sweet potatoes, 1119
DDT, conversion to DDE, 528
DDT on peach in Ontario, 109
DDT, trend of on apples, 1053
demeton, on various vegetables, 741
Dibrom, 353
Dieldrin in cattle, 1206
Dieldrin on and in lemons, 42
endrin, spray coverage, 592
guthion, sampling for, 791
heptachlor in cattle, 1026
heptachlor on and in lemons, 42
Hercules AC-528, mechanism of loss of, 111
Hercules AC-528 in milk, 349
insecticides, for
control of cheese mites, 237
control of dermestids, 234
control of imported fire ant, 1
control of Japanese beetle, 9
protection of stored woolens, 457
lindane, in sheep and goats, 1031
malathion, on and in oranges, 42
methoxychlor, in milk, 341, 866
methoxychlor on vegetable crops, 232
organophosphorus insecticides, 840
parathion, trend of on apples, 1053
phosdrin, on pin oak, 1015
pyrethrins, on wheat, 703
Tedion, location of on citrus fruit, 723
Thimet, 304, 353
Thiodan, on corn plants, 102
Thiodan, effect of grain moisture content on, 131
toxaphene, in cattle, 1206
Resistance of
alfalfa to spotted alfalfa aphid, 949
cattle to cattle grubs, 1022
boll weevil to insecticides, 555
brown dog tick to lindane, chlordane, 332
corn leaf tissue to European corn borer, 1171
Culicoides furens to various compounds, 352
Diazinone-resistant house flies to insecticides, 733
Dilan-resistant house flies to Prolan and Bulan, 997
European house flies to insecticides, 419
German cockroach to insecticides, 544
house flies and X_5 ratio, 1077
Malathion-resistant flies to carbamates, 178
Orlando house fly colonies to carbamates, 308
pear psylla to dieldrin, 530
prevention of by spray rotation, 432
small grains to greenbug, 276
spider mites to insecticides, 936
Tetranychus cinnabarinus to organic phosphates, 760
in two-spotted spider mite, 1026
Reticulitermes flavipes, 533
Rhabditis spp., 861
Rhagoletis
cingulata indifferens, 769
completa, 279
pomonella, 751
Rhipicephalus sanguineus, 332
Rhizoglyphus echinopus, 188
Rhodnius prolixus, 1096
Rhopalosiphoninus latysiphon, 424
Rhopalosiphum
fitchii, 8, 422, 423, 621, 994
junackianum, 424
maidis, 8, 424, 440, 972
nymphaeae, 8, 422, 424
pseudobrassicae, 8
rufomaculatum, 424
tanacetina, 424
Rhyacionia
buoliana, 588, 768
frustrana, 1180
"*Rhynchagrotis*"
exsertistigma, 966
insularis, 966
Rhyzopertha dominica, 557, 888
Rohm-Haas C-140, for control of
apple aphid, 493
rosy apple aphid, 493
Rodolia cardinalis, 72, 1099
Ronnel (see also Korlan, Trolene, Dow-ET-57, Dow ET-14) effect of on cattle grubs and cattle weight gain, 488
fathead minnow, 331
garden symphylid, 672
goat lice, 980
horn flies, 1216
house flies, 704, 779
Psila rosae, 963
screw-worm, 1217
Ronnell, for control of
cattle grubs, 524, 958, 1028, 1222
Drosophila melanogaster, 1136
Drosophila spp., 755
horn fly, 648
potato aphid, 355
spotted alfalfa aphid, 714
Tendipes plumosus, 331
tomato suckfly, 1219
winter tick, 1220
Ronnell, toxicity of to house flies, 733
Rotenone, effect of on Japanese beetle, 9
Rotenone, for control of
alfalfa weevil, 663
beet leafhopper, 701
cabbage caterpillars, 221
horn flies, 762
meadow spittlebug, 663
pea aphid, 663
sheep ked, 380
Rungsia maydis, 424
Ryania, effect of on
apple insects, 819
sugarcane borer, 821
Ryania, for control of corn earworm, 1111

Sabadilla, effect of on harlequin cabbage bug, 777
Saissetia
hemisphaerica, 596
nigra, 596
oleae, 74, 596
Salticus scenicus, 530
Sampling of
alfalfa for estimation of Guthion residues, 791
red-pine sawfly, 600
spruce budworm, 618
Sanninoidea exitiosa graefi, 804
Santocel C, effect of on poultry ectoparasites, 1195
Sassafras oil, tests of repellency of, 286
Scambus
alboricta, 550
aplopappi, 531
hispae, 550
Scatophaga stercoraria, 530
Scelio pembertoni, 73
Scheloribates sp., 188
Schradan, effect of soils on uptake of, 1160
Schradan, effect of on
alfalfa mites, 991
garden symphylid, 672
Schradan, for control of
citrus whitefly, 1008
cotton thrips, 35
European red mite, 936
green peach aphid, 727
two-spotted spider mite, 1026
Sciopithes obscurus, 3
Scolia ruficornis, 71
Scolothrips sexmaculatus, 936
Scolytus
multistriatus, 902
ventralis, 323
Scotogramma trifolii, 1212
Scutellista cyanea, 596
Scutigerella immaculata, 666, 672, 678, 892
Scymnus flavifrons, 596
SD-51, for control of cattle grubs, 1222
SD-75, for control of cattle grubs, 1222
Seeds, germination of
effect of insecticides on, 355, 360
effect of insecticides on
barley, 319
oats, 415
rice, 706
wheat, 51, 535, 650
Seeds, treatment of, for control of
beet leafhopper, 700
cotton pests, 807
rice water weevil, 767
spotted alfalfa aphid, 654, 714
wireworms, 509
Septis
arctica, 966
cuculliformis, 966
finitima, 966
dl, Serine, for control of beet leafhopper, 701
Sevin
bioassay of residues of, 840
effectiveness of on vegetable crops, 477
resistance to of pear psylla, 530

systemic action in control of northern fowl mite, 170
Sevin, effect of on
alfalfa weevil, 758
apple insects, 826
cotton insects, 826
grey garden slug, 360
harlequin cabbage bug, 777
honey bee, 549
horn flies, 1216
insects attacking Douglas-fir logs, 482
Japanese beetle, 9
pea aphid, 758
potato leafhopper, 908
poultry ectoparasites, 1195
Psila rosae, 963
sugarcane borer, 821
Sevin, for control of
alfalfa weevil, 758
apple aphid and rosy apple aphid, 493
beet leafhopper, 701
corn earworm, 338, 1111
European red mite, 936
harlequin cabbage bug, 779
lesser peach twig borer, 637
meadow spittlebug, 663
pea aphid, 663
potato aphid, 356
thrips on spinach, 540
tobacco hornworm, 798
winter ticks, 1220
Sevin, toxicity of to
Douglas-fir beetle, 485
house flies, 733
plum curculio, 921
Shell OS-1808, effect of on alfalfa mites, 991
Shell 3562, for control of corn earworm, 1111
Shell SD 4402, for control of
Conoderus falli, 780
corn earworm, 1111
tobacco hornworm, 798
Sigmodon hispidus, 770
Simulium
decorum, 629
fibrinflatum, 629
jenningsi, 629
parnassum, 629
tuberosum, 629
venustum, 629
vittatum, 629
Sinea confusa, 159
Sinella curviseta, 824
Sipha agropyrella, 424
Siphona irritans, 341, 648, 762, 866, 1216
Sitodiplosis mosellana, 72
Sitona
flavescens, 1155
hispidula, 1155
Sitophilus
granarius, 131, 443, 557, 621
oryza, 443, 557, 612, 621, 888, 942, 1091, 1130
Sitotroga cerealella, 557, 612, 621, 888
Sodium chloride, effect of on *Acarus siro*, 237
Soils, effect of on
chemical control of onion maggot, 851
uptake of systemics by plants, 1160
Solenopsis
molesta, 365
saevissima var. *richteri*, 1, 347, 455
Solenotus
begini, 690, 898
pulchripes, 898
websteri, 898
Sorbic acid, for control of *Acarus siro*, 237
Sparganothus xanthoides, 836
Spergon (see chloronil)
Spilochalcis side, 531
Spilonotus ocellana, 1103
Spissistilus festinus, 286, 429
Sprays, for control of
cotton insects, 466, 985
house flies, 704
water rice weevil, 767
Sprays
conventional vs. high-clearance, for cotton, 592
logarithmic, for screening insecticides, 534
Spread of Japanese beetle, 771
Staining of symbionts of confused flour beetle, 756
Stains, attraction of to fabric pests, 382
Starvation as cause of defective honey bee eggs, 166
Stegasta basqueella, 468
Stephanitis
globulifera, 712
pyrioides, 712
Sterile males, release of screwworm in Florida, 1202
Sterility, induced in
Anopheles quadrimaculatus by radiation, 868
Tetranychus telarius by Tedion, 535
Stethorus
atomus, 229
punctum, 819
spp., 936
Stimulation, electrical, release of neuroactive agent by in American cockroach, 1070
Stomoxys calcitrans, 866
Storage rack for vials, 1225
Strobane, for control of
Acarus siro, 237
alfalfa weevil, 663
larger cabinet beetle, 234
meadow spittlebug, 663
pea aphid, 663
Tyroglyphus spp., 237
Sucrose, attractants for house flies in, 981
Sugars secreted by woolly alder aphid, 353
Sulfur, wettable, effect of on cotton pests, 975
Entomophthora spp., 28
Sulfur, wettable, for control of cabbage looper, 1013
Surveys of Japanese beetle, 771
Swammerdamia spp., 1099
Symbionts in confused flour beetle, absence of, 756
Sympiesis sp., 531
Synanthedon pictipes, 634
Syneches sp., 430
Synergists, analog of DDT as synergists of DDT, 626
Systemics
Dowco 109 as systemic for animals, 749
systemic properties of Bayer 25141 and analogues, 99
Systemics, for control of
cabbage aphid, 1045
cattle grubs, 1004, 1022, 1028, 1222
grasshopper damage, 37
house fly larvae, administered to chicks, 446, 719
Myzus cerasi, 778
Tabanus spp., 884
Tarsonemus confusus, 991
TDE (see also DDD)
effect of on seed germination, 355
inducing sterility of *Tetranychus telarius*, 535
locale of residues of on citrus fruits, 723
TDE, effect of on
apple insects, 826
Japanese beetle, 9
lepidopterous larvae, 1115
TDE, for control of
army cutworm, 785
tobacco hornworm, 798
TDE, toxicity of to house flies, 733
Tedion, effect of on poultry ectoparasites, 1195
Tedion, for control of
European red mite, 936
mites in insect cultures, 443
two-spotted spider mite, 342, 1026
Telenomus spp., 895
Telone, for control of garden symphylid, 678
10-undecenoic acid, repellency of to pea aphid, 106
Tendipes plumosus, 331
Tenebroides mauritanicus, 557, 612
Temperature, effect of on spotted alfalfa aphid resistance in alfalfa, 949
Tephrosia, for control of
alfalfa weevil, 663
meadow spittlebug, 663
pea aphid, 663
TEPP
effect of on fathead minnows, 331
for control of *Tendipes plumosus*, 331
penetration of to cockroach nerve cord, 812
Tetrachloroethane, for control of western peach tree borer, 804
Tetrachlorophthalic acid, for control of beet leafhopper, 701
Tetram,
effect of on fathead minnows, 331
for control of *Tendipes plumosus*, 331
penetration of to cockroach nerve cord, 812
quaternarized, penetration of to cockroach nerve cord, 812
Tetranychus
atlanticus, 188
canadensis, 188
cinnabarensis, 188
cinnabarinus, 279, 760, 807
desertorum, 188, 757, 975
hicoriae, 163
marianae, 763
mcdanieli, 188
pacificus, 936
telarius, 189, 342, 411, 461, 477, 535, 807, 871, 991, 1015, 1026
tumidus, 189
Tetraleurodes
acaciae, 9
mori, 9
Tetraticia sp., 9
Tetrastichus bruchophagi, 898
Thanite, effect of on biting flies of cattle, 866
Thaumatomya glabra, 430
Thelazia
californiensis, 1030
rhodesii, 1030
Therioaphis maculata, 8, 28, 136, 142, 154, 249, 368, 430, 431, 440, 537, 642, 654, 714, 949, 1023, 1057, 1099

Thermobia domestica, 448
Theronia atalantae, 550
Thimet (see also phorate)
effect of on germination and seedling growth of wheat, 650
effect of on rate of seeding wheat, 535
preassay activation of, 353
for prevention of grasshopper damage, 37
Thimet, effect of on
cotton and cotton insects, 807
cotton aphid, 304
fathead minnows, 331
garden symphylid, 666, 672
greenbug, 304
harlequin cabbage bug, 777
Japanese beetle, 9
Psila rosae, 963
Thimet, for control of
alfalfa weevil, 663
andromeda lace bug, 712
apple aphid, 493
azalea lace bug, 712
birch leaf miner, 712
boxwood leaf miner, 712
bronze birch borer, 255
cabbage aphid, 1045
citrus whitefly, 1008
Conoderus falli, 780
cotton thrips, 35
Dasyneura gleditschiae, 712
Drosophila spp., 755
flat headed apple tree borer, 255
green peach apid, 727
holly leaf miner, 712
meadow spittlebug, 663
Myzus cerasi, 778
pea aphid, 663, 1212
rosy apple aphid, 493
spotted alfalfa aphid, 654, 714
Tendipes plumosus, 331
tobacco thrips, 451
Thimet, toxicity of to
European pine shoot moth, 588
onion maggot, 683
Thiodan, effect of on
alfalfa weevil, 758
cabbage aphid, 816
cabbage looper, 816
chinch bug, 759
corn earworm, 338
Estigmene acrea, 1109
harlequin cabbage bug, 777
insects attacking Douglas-fir logs, 482
Japanese beetle, 9
lesser cornstalk borer, 63
Lygus bug, 461
Myzus persicae, 781
pea aphid, 758
Pieries rapae, 816
potato insects, 564
potato leafhopper, 908
Psila rosae, 963
sugarcane borer, 821
Thiodan, for control of
alfalfa weevil, 663
apple aphid, 493
army cutworm, 785
beet heafhopper, 701
birdsfoot trefoil insects, 836
cabbage looper, 1013
corn earworm, 338, 1111
cotton insects, 1068
European corn borer, 102
lesser peach twig borer, 637
meadow spittlebug, 534, 663
Mexican bean beetle, 541
pea aphid, 663
rosy apple aphid, 493
thrips on spinach, 540
tobacco hornworm, 798
tomato suckfly, 1219
western peach tree borer, 804
Thiodan, toxicity of to
Douglas-fir beetle, 485
western cherry fruit fly, 769
Thiram
effect of on apple insects, 826
for control of onion maggot, 851
Thrips tabaci, 807
Thyridopteryx ephemeraeformis, 353
Thysanus flavopalliatus, 577
Tineola bisselliella, 382, 457
Tiphia
vernalis, 71
spp., 1099
Tolerance of potato flea beetle to DDT and dieldrin, seasonal, 519
Toxaphene,
action of on resistant bollweevils, 403
bioassay of residues of, 840
effect of on red clover yields, 1145
residues of on sweet potato, 1119
resistance of pear psylla to, 530
Toxaphene, effect of on
alfalfa arthropods, 428
alfalfa mites, 991
beet armyworm, 877
boll weevil, 1042
cabbage aphid, 816
cabbage looper, 816, 877
chinch bug, 759
cotton, 807
cotton insects, 807, 975, 985
Estigmene acrea, 1109
goat lice, 980
harlequin cabbage bug, 777
Japanese beetle, 9
lepidopterous larvae, 1115
Lygus bugs, 461
pecan aphids and mites, 163
Pieris rapae, 816, 877
potato leafhopper,908
sugarcane borer, 821
Toxaphene, for control of
alfalfa weevil, 663, 939
army cutworm, 875
cabbage caterpillars, 221
cabbage looper, 1013
corn earworm, 1111
cotton insects, 1068
cotton thrips, 35
lesser peach twig borer, 637
meadow spittlebug, 663
pea aphid, 663
Pyrausta nubilalis, 49
sheep ked, 380
Stegasta basqueella, 468
tobacco thrips, 451
winter tick, 1220
Toxaphene, residues of in cattle, 1206
Toxaphene, toxicity of to
convergent lady beetle, 616
onion maggot, 683
Orius insidiosus, 616
pink bollworm, 301
western cherry fruit fly, 769
Toxoptera graminum, 8, 276, 304, 368, 536
Tractors, use of in applying insecticides, 1016
Traps
aromatic bait, as factor in insect spread, 157
baffle designs in, 167
blacklight, for European chafer, 744
blacklight, for various insects, 1212
measurement of air volume displacement, 343
method of testing of, 496
pollen, for honey bee hives, 1024
Traps, for
Culex tarsalis, studies on, 496
Hippelates pusio, studies on, 496
insects associated with water surface, 348
Trialeurodes
abutilonea, 9
floridensis, 9
Tribolium
castaneum, 612, 888, 1184
confusum, 99, 618, 756, 942, 1081, 1091, 1184
sp., 557
Trichogramma
minutum, 550, 768
spp., 1099
Trichoplusia ni, 221, 447, 816, 877, 1013, 1045, 1068
Trimeromicrus maculatus, 898
Trimerotropus pallidipennis, 337
Triozys
utilis, 136, 142, 154
sp., 536
Trioza collaris, 9
Trithion, effect of on
alfalfa mites, 991
cabbage aphid, 816
cabbage looper, 816
chinch bug, 759
Entomophthora spp., 28
fathead minnow, 331
garden symphylid, 666, 672
harlequin cabbage bug, 777
Japanese beetle, 9
Trithion, for control of
alfalfa weevil, 590, 663
apple aphid, 493
beet leafhopper, 700
European red mite, 936
lesser peach twig borer, 637
meadow spittlebug, 663
pea aphid, 663
potato aphid, 355
rosy apple aphid, 493
scale insects, on dormant fruit trees, 373
spotted alfalfa aphid, 654
Tendipes plumosus, 331
Tetranychus cinnabarinus, 760
Tetranychus marianae, 763
thrips on spinach, 540
tobacco hornworm, 798
tomato suckfly, 1219
Trogoderma
glabrum, 830
granarium, 51, 312, 830
grassmani, 830
inclusum, 234, 830
ornatum, 830
parabile, 830
simplex, 830
sternale, 830
Trolene (see also ronnel) for control of cattle grubs, 1022
Trombicula akamushi, 824
Tropic acid esters, toxicity of to house flies, 954
Tydeidae, 725
Tydeus sp., 189, 231, 991
Tylencholaimellus striatus, 861
Tylenchorhynchus maximus, 861

Tylenchus
costatus, 861
davainii, 861
exiguus, 861
Typhlodromus
conspicuus, 231
cucumeris, 991
fallacis, 871
floridanus, 231
perigrinus, 577
sp., 190, 231, 443, 819, 871, 936
Tyroglyphus castellanii, 514
Tyrophagus sp., 189, 237

UCC-8305
effect of on garden symphylid, 672
effect of on *Psila rosae*, 963
for control of corn earworm, 1111
Uranotaenia sapphirina, 1050

Vanessa cardui, 1017
Vanside F-1562, for control of
Acarus siro, 237
larger cabinet beetle, 234
Vapam, for control of
garden symphylid, 678
strawberry weevils, 3
Vapors of lindane and aldrin to control adult mushroom flies, 257
Vasates sp., 189
VC-13, effect of on
garden symphylid, 666, 672
fathead minnows, 331
Psila rosae, 963
VC-13, for control of *Tendipes plumosus*, 331
Veratraldehyde, acetal, toxicity of to house flies, 954
Viruses
beet yellows, 910
cucumber mosaic, 421
curly top, 470, 700
feathery mottle virus, 1057
lettuce yellows, 686
potato leaf roll, 727
Tristeza (T_2 severe strain), 632
Viruses, granulosis of
Alaskan insects, 350
Recurvaria milleri, 531

Weather in control of insect populations, 532
Weight
of queen honey bees controlled genetically, 547
relationship of to effect of insecticides, 1115
White oil, for control of citrus whitefly, 1008
Wohlman salts
effect of on drywood termite, 1211
tests of with *Reticulitermes flavipes*, 533

Xenopsylla cheopis, 770
Xiphenema
americanum, 861
diversicaudatum, 861
Xylena
nupera, 966
spp., 1103
Xyletinus peltatus, 452

Zagrammosoma
americanum, 531
mirium, 895
nigrolineatum, 690
spp., 895
Zelus renardii, 159
Zercoseius sp., 577
Zinc chloride, effect of on drywood termite, 1211
Zinc sulfate compatibility of with Thiodan on potato foliage, 564
Zosteropoda hirtipes, 966